

CERN-TH/2002-330

IFT - 40/2002

hep-ph/0211371

November 2002

TWO-HIGGS-DOUBLET MODELS WITH CP VIOLATION *

I. F. Ginzburg¹, M. Krawczyk^{2,3} and P. Osland⁴

¹ *Sobolev Institute of Mathematics, SB RAS, 630090 Novosibirsk, Russia*

² *Theory Division CERN, CH-1211 Geneva 23, Switzerland*

³ *Institute of Theoretical Physics, Warsaw University, Poland*

⁴ *Department of Physics, University of Bergen, N-5007 Bergen, Norway*

Abstract

We consider the Two-Higgs-Doublet Model and determine the range of parameters for which CP violation and Flavor Changing Neutral Current effects are naturally small. It corresponds to small values of the mass parameter m_{12}^2 , describing soft (ϕ_1, ϕ_2) mixing in the potential. We discuss how, in this approach, some Higgs bosons can be heavy, with mass of the order of 1 TeV.

The possibility that at the Tevatron, LHC and an e^+e^- Linear Collider, only one Higgs boson will be found, with properties indistinguishable from those in the Standard Model (SM), we define as the SM-like scenario. While this scenario can be obtained with large $\mu^2 \sim \text{Re } m_{12}^2$ parameter, in which case there is decoupling, we here discuss the opposite case of small μ^2 , without decoupling.

CERN-TH/2002-330

IFT - 40/2002

hep-ph/0211371

November 2002

*Presented at SUSY02, DESY, June 2002 and LCWS02, Jeju, Korea, August 2002

1 Introduction

We consider the following Two-Higgs-Doublet Model (2HDM) potential, with quartic and quadratic terms separated [1, 2, 3, 4, 5]:

$$\begin{aligned}
V = & \frac{1}{2}\lambda_1(\phi_1^\dagger\phi_1)^2 + \frac{1}{2}\lambda_2(\phi_2^\dagger\phi_2)^2 + \lambda_3(\phi_1^\dagger\phi_1)(\phi_2^\dagger\phi_2) + \lambda_4(\phi_1^\dagger\phi_2)(\phi_2^\dagger\phi_1) \\
& + \frac{1}{2}[\lambda_5(\phi_1^\dagger\phi_2)^2 + \text{h.c.}] + \{[\lambda_6(\phi_1^\dagger\phi_1) + \lambda_7(\phi_2^\dagger\phi_2)](\phi_1^\dagger\phi_2) + \text{h.c.}\} \\
& - \{m_{11}^2(\phi_1^\dagger\phi_1) + [m_{12}^2(\phi_1^\dagger\phi_2) + \text{h.c.}] + m_{22}^2(\phi_2^\dagger\phi_2)\}.
\end{aligned} \tag{1}$$

As is well known, both CP violation in the Higgs sector and flavor-changing neutral currents (FCNC) can be suppressed by imposing a Z_2 symmetry [6]. This requires symmetry of the potential under $(\phi_1 \rightarrow -\phi_1, \phi_2 \rightarrow \phi_2)$ (or vice versa), which implies $\lambda_6 = \lambda_7 = m_{12}^2 = 0$. We shall allow *soft* violation of this symmetry, i.e., we take $\lambda_6 = \lambda_7 = 0$, but allow $m_{12}^2 \neq 0$ [3, 5, 7]. A simple discussion can be given for this case, in which $\text{Im } m_{12}^2 \neq 0$ signals CP violation.

2 CP violation

We shall now consider the simpler case of $\lambda_6 = \lambda_7 = 0$, and parametrize the minimum of the potential (or vacuum) as

$$\phi_1 = \begin{bmatrix} 0 \\ \frac{1}{\sqrt{2}}v_1 \end{bmatrix}, \quad \phi_2 = \begin{bmatrix} 0 \\ \frac{1}{\sqrt{2}}v_2 e^{i\xi} \end{bmatrix}. \tag{2}$$

Naively, the phase ξ violates CP, but it can be removed by a global phase transformation on the field ϕ_2 , together with the phases of λ_5 , m_{12}^2 and the fermion fields [3, 8]. It is convenient to define

$$\mu^2 = \text{Re}(m_{12}^2 e^{i\xi}) \frac{v^2}{v_1 v_2}. \tag{3}$$

The phase ξ can be found from the equation

$$\text{Im}(m_{12}^2 e^{i\xi}) = \text{Im}(\lambda_5 e^{2i\xi})v_1 v_2. \tag{4}$$

Making use of the rephasing invariance [3, 8], we put $\xi = 0$. With this choice, eq. (4) becomes a constraint for the relation of $\text{Im}(m_{12}^2)$ to $\text{Im } \lambda_5$.

The neutral sector has a mass squared matrix of the form

$$\mathcal{M}^2 = \begin{pmatrix} \mathcal{M}_{11}^2 & \mathcal{M}_{12}^2 & -\frac{1}{2}\text{Im } \lambda_5 v^2 \sin \beta \\ \mathcal{M}_{12}^2 & \mathcal{M}_{22}^2 & -\frac{1}{2}\text{Im } \lambda_5 v^2 \cos \beta \\ -\frac{1}{2}\text{Im } \lambda_5 v^2 \sin \beta & -\frac{1}{2}\text{Im } \lambda_5 v^2 \cos \beta & \mathcal{M}_{33}^2 \end{pmatrix} \tag{5}$$

where $\mathcal{M}_{11}^2$, $\mathcal{M}_{12}^2$, $\mathcal{M}_{22}^2$ and $\mathcal{M}_{33}^2$ are the same as in the CP-conserving case. When $\text{Im } \lambda_5 = 0$, there is no CP violation, the matrix (5) is block diagonal, and the physical states are h , H and A . When $\text{Im } \lambda_5 \neq 0$, all three neutral Higgs states mix; we denote them by h_1 , h_2 and h_3 .

The mass-squared matrix may be diagonalized via a rotation matrix, defined by

$$R \mathcal{M}^2 R^T = \text{diag}(M_1^2, M_2^2, M_3^2). \quad (6)$$

In the limit of weak CP violation, the masses for h_1 , h_2 and h_3 will deviate from those of h , H and A by terms quadratic in $\text{Im } \lambda_5$.

3 Decoupling or no decoupling?

We shall here consider the scenario of weak (or no) CP violation and large masses of Higgs particles except one, namely $M_{h_1} \sim M_h$. Let us discuss how large masses M_A (close to M_{h_3}) and $M_{H^\pm}$ arise in such a case. The potential (1) (but with $\lambda_6 = \lambda_7 = 0$) gives

$$M_A^2 = \frac{1}{2}\mu^2 - \text{Re } \lambda_5 v^2 \quad \text{and} \quad M_{H^\pm}^2 = \frac{1}{2}[\mu^2 - (\lambda_4 + \text{Re } \lambda_5)v^2]. \quad (7)$$

There are two rather distinct mechanisms for obtaining large mass M_A^2 and $M_{H^\pm}^2$: either (i) μ^2 is large (this is extensively discussed by Haber as *the decoupling scenario*) [2, 5], or (ii) μ^2 is small, whereas $|\text{Re } \lambda_5|$ is “large” [4, 5]. In the latter case, there are obvious upper bounds (from perturbativity and positivity) on how large $|\text{Re } \lambda_5|$ can be. Decoupling properties of the Two-Higgs-Doublet Model were studied in [9].

In this model, with weak (or no) CP violation, one can realize a Standard-Model-Like Scenario:

- *There is a light Higgs boson with couplings to the up (e.g. t) and down (e.g. b) type quarks, and to W and Z , like in the Standard Model,*

$$|g_i| \approx |g_i^{\text{SM}}| \quad (i = W, Z, \text{down}, \text{up}). \quad (8)$$

- *The other Higgs bosons are heavy, $\mathcal{O}(1 \text{ TeV})$.*

Within the Two-Higgs-Doublet Model, this scenario can be realized in two distinct ways. They are [4]:

- Solutions A. All basic couplings are approximately the same as in the SM, up to an overall sign.
- Solutions B. Like Solutions A, *except that the couplings to either up- or down-type quarks have opposite signs of those in the SM.* This case cannot be realized in the decoupling scenario.

4 Model II. Observables

Let us now be more specific, and consider the so-called Model II for Yukawa couplings, where masses of down- and up-type quarks originate from couplings to ϕ_1 and ϕ_2 , respectively. We denote by $\chi_V^{h_i}$, $\chi_u^{h_i}$ and $\chi_d^{h_i}$ the ratios of the Higgs couplings to W and Z (V) and to up and down-type quarks, with respect to those of the Standard Model. In particular, for the Yukawa couplings these ratios can be expressed via elements of the rotational matrix R of eq. (6) as

$$\chi_u^{h_i} = \frac{1}{\sin \beta} [R_{i2} - i\gamma_5 \cos \beta R_{i3}], \quad \chi_d^{h_i} = \frac{1}{\cos \beta} [R_{i1} - i\gamma_5 \sin \beta R_{i3}], \quad (9)$$

where R_{i3} is proportional to $\text{Im } \lambda_5$. Note that in accordance with eq. (9), the CP violation induced by Higgs exchange in $t\bar{t}$ production [10] provides information on $\text{Im } \lambda_5$.

Furthermore, these relative couplings satisfy a *pattern relation* [4, 8]:

$$(\chi_u^{h_i} + \chi_d^{h_i})\chi_V^{h_i} = 1 + \chi_u^{h_i}\chi_d^{h_i}. \quad (10)$$

In the CP -conserving case, even with all basic couplings being the same (up to a sign) as in the SM (8), loop-induced transition rates, like $h \rightarrow \gamma\gamma$, may differ from the SM prediction. This is due to the different behaviors of the trilinear Higgs coupling hH^+H^- for small and large μ . In fact, the ratio of this coupling to its SM value can be written as

$$\chi_{H^\pm}^h \equiv -\frac{vg_{hH^+H^-}}{2M_{H^\pm}^2} = \left(1 - \frac{M_h^2}{2M_{H^\pm}^2}\right)\chi_V^h + \frac{M_h^2 - \frac{1}{2}\mu^2}{2M_{H^\pm}^2}(\chi_u^h + \chi_d^h). \quad (11)$$

Thus, if $\mu^2 \sim M_{H^\pm}^2$, there is no effect in $\Gamma_{\gamma\gamma}$, whereas if $\mu^2 < M_{H^\pm}^2$ there is a difference of several per cent, as illustrated in Fig. 1 (left panel) for the case of Solutions A [4]. Non-decoupling effects in the 2HDM were studied for other processes in [11, 12].

These deviations from unity are large enough that the form of the 2HDM potential (large or small μ) can be tested at a $\gamma\gamma$ Collider [13].

Also, the loop-induced couplings to two gluons may differ from those of the SM-value, but this occurs only for Solutions B. This effect is also illustrated in Fig. 1 (right panel).

Acknowledgements

We are grateful to the CERN, TH Division for kind hospitality and support during work on this paper. We thank Maria Jose Herrero Solans for informing us about ref. [9]. I.G. is also grateful to the Landau–Centro Volta Network and to Prof. O. Panella for the warm hospitality during a stay at INFN Perugia when the final version of this paper was prepared. This work is supported in part by RFBR grants 02-02-17884 and 00-15-96691 and INTAS 00-00679, by the Polish Committee for Scientific Research (grant No. 5 P03B12120 (2002)) and by the European Community's Human Potential Programme under contract HPRN-CT-2000-00149 Physics at Colliders, and by the Research Council of Norway.




Figure 1: Ratios of the Higgs boson decay widths in the SM-like 2HDM (II) and the SM as functions of M_h . *Left panel:* $h \rightarrow \gamma\gamma$ decay widths, solutions A, for $M_{H^\pm} = 800$ GeV and $\mu/\sqrt{2} = x M_{H^\pm}$. *Right panel:* $h \rightarrow gg$, solutions B.

References

- [1] J. L. Diaz-Cruz and A. Mendez, Nucl. Phys. B**380** (1992) 39,
J. L. Diaz-Cruz and G. Lopez Castro, Phys. Lett. B**301** (1993) 405.
- [2] H. E. Haber, arXiv:hep-ph/9501320.
- [3] G. C. Branco, L. Lavoura, J. P. Silva, *CP Violation* (Oxford Univ. Press, 1999)
- [4] I. F. Ginzburg, M. Krawczyk and P. Osland, Linear Collider Note LC-TH-2001-026, In *2nd ECFA/DESY Study 1998–2001*, 1705 [arXiv:hep-ph/0101208];
I. F. Ginzburg, M. Krawczyk and P. Osland, Nucl. Instrum. Meth. A**472** (2001) 149 [arXiv:hep-ph/0101229].
- [5] J. F. Gunion and H. E. Haber, arXiv:hep-ph/0207010.
- [6] S. L. Glashow and S. Weinberg, Phys. Rev. D**15** (1977) 1958.
- [7] R. Santos, S. M. Oliveira and A. Barroso, arXiv:hep-ph/0112202.
- [8] I. F. Ginzburg, M. Krawczyk and P. Osland, to be published.
- [9] P. Ciafaloni and D. Espriu, Phys. Rev. D **56** (1997) 1752 [arXiv:hep-ph/9612383].
- [10] W. Bernreuther and A. Brandenburg, Phys. Rev. D**49** (1994) 4481 [arXiv:hep-ph/9312210].
- [11] M. Malinsky, arXiv:hep-ph/0207066.

- [12] S. Kanemura, S. Kiyoura, Y. Okada, E. Senaha and C. P. Yuan, arXiv:hep-ph/0209326; arXiv:hep-ph/0211308.
- [13] J. A. Aguilar-Saavedra *et al.* [ECFA/DESY LC Physics Working Group Collaboration], “TESLA Technical Design Report Part III: Physics at an e+e- Linear Collider,” arXiv:hep-ph/0106315.